

A proof of Björner's totally nonnegative conjecture

Michael Björklund Alexander Engström*

February 2, 2008

Abstract

The McMullen Correspondence gives a linear dependence between M-sequences of length $\lfloor \frac{d}{2} \rfloor + 1$ and f -vectors of simplicial d -polytopes. Denote the transfer matrix between g and f by M_d . Recently, Björner proved that any 2×2 -minor of M_d is nonnegative and conjectured that the same would be true for arbitrary minors. In this paper we answer the question in the affirmative.

1 Introduction

Let P be a d -dimensional simplicial polytope, and let f_i denote the number of i -dimensional faces. The nonnegative integer vector

$$f = (f_0, \dots, f_{d-1})$$

is called the f -vector of P . We assume the convention $f_{-1} = f_d = 1$. The Euler-Poincaré formula gives a first rough estimate on the dependence between the elements of f ,

$$-f_{-1} + f_0 - f_1 + \dots + (-1)^{d-1} f_{d-1} + (-1)^d f_d = 0.$$

Define the numbers $g_k = h_k - h_{k-1}$ for $k = 0, \dots, \lfloor \frac{d}{2} \rfloor$, where

$$h_i = \sum_{j=0}^i (-1)^{i+j} \binom{d-j}{i-j} f_{j-1}, \quad i = 0, \dots, d.$$

The nonnegative vector $g = (g_0, \dots, g_{\lfloor \frac{d}{2} \rfloor})$ will be referred to as the g -vector of the simplicial d -polytope P . The McMullen correspondence asserts that the map

$$g \mapsto g \cdot M_d,$$

where M_d is a nonnegative $(\lfloor \frac{d}{2} \rfloor + 1) \times d$ matrix given by

$$m_{ij} = \binom{d+1-i}{d-j} - \binom{i}{d-j} \quad i = 0, \dots, \lfloor \frac{d}{2} \rfloor \quad \text{and} \quad j = 0, \dots, d-1,$$

*{mickebj,alexe}@math.kth.se, Department of Mathematics, Royal Institute of Technology, S-100 44 Stockholm, Sweden.

is a bijection between M -sequences g with $g_1 = n - d - 1$ and f -vectors in $\mathbb{N}_0^d$ of simplicial d -polytopes with $n = g_1 + d + 1$ vertices. Recall that a sequence $n_0, n_1, \dots$, of nonnegative integers is a M -sequence if $n_0 = 1$ and

$$\partial^k(n_k) \leq n_{k-1} \quad \text{for all } k > 1,$$

where ∂^k is the “ k -boundary”-operator (see e.g. [8], 262). A theorem by Macaulay [6] gives an algebro-combinatorial characterization of such sequences.

This paper is devoted to the analysis of the transfer matrices M_d above. In dimensions $d = 1, 2$, and 3, they are given by

$$M_1 = \begin{pmatrix} 1 & 2 \end{pmatrix} \quad M_2 = \begin{pmatrix} 1 & 3 & 3 \\ 0 & 1 & 1 \end{pmatrix} \quad M_3 = \begin{pmatrix} 1 & 4 & 6 & 4 \\ 0 & 1 & 3 & 2 \end{pmatrix}.$$

Björner proved in [2] that every 2×2 -minor in M_d is nonnegative. In the same paper he conjectured that any minor in M_d is nonnegative (such a matrix is called *totally nonnegative*). The conjecture has been verified up to dimension $d = 13$ by A. Hultman. In this paper we settle the conjecture in the affirmative for all dimensions.

2 Lattice paths and nonnegative minors

A path from (x_1, y_1) to (x_2, y_2) in $\mathbb{Z}^2$, where $x_1 \leq x_2$ and $y_1 \leq y_2$, is called a lattice path if only the steps $(1, 0)$ and $(0, 1)$ are allowed. The number of lattice paths from $(0, 0)$ to (m, n) which do not touch the line $y = x + t$ are $\binom{m+n}{n} - \binom{m+n}{m-t}$ if $t > 0$ [1, 7]. This is sometimes referred to as the ballot numbers. The weight of a path is the product of the weights of its arcs. From any subset A of $\mathbb{Z}^2$ one can construct a planar acyclic directed graph with vertex set A and arcs of types $(x, y) \rightarrow (x+1, y)$ and $(x, y) \rightarrow (x, y+1)$.

Theorem 2.1 *The matrix M_d is totally nonnegative for all d .*

PROOF: For integers $n \geq 2$ define the graphs T_n with vertex set

$$\{(x, y) \in \mathbb{Z}^2 \mid x \leq \lceil n/2 \rceil - 1, y - x \leq \lfloor n/2 \rfloor, \text{ and } x + y \geq \lceil n/2 \rceil - 1\}.$$

The weight of horizontal arcs in T_n is 1, and the weight of any vertical arc $(x, y) \rightarrow (x, y+1)$ is w_y . The graphs T_8 and T_9 are depicted in figures 1 and 2.

For all $0 \leq i \leq \lceil n/2 \rceil - 1$ and $0 \leq j < n$, let the sum of the weights of the directed paths from $(\lceil n/2 \rceil - 1 - i, i)$ to $(\lceil n/2 \rceil - 1, j)$ be $W(n, i, j)$. If $i > j$ then $W(n, i, j) = 0$ and otherwise it is

$$w_i w_{i+1} \cdots w_{j-1} \left(\binom{j}{i} - \binom{j}{n-i} \right)$$

by the ballot numbers.




Figure 1: The weighted planar graph T_8




Figure 2: The weighted planar graph T_9

Now define the weights of the vertical arcs as $w_i = \binom{n}{i+1} / \binom{n}{i}$. Note that all arc weights are positive real numbers. For $i \leq j$ we get that

$$\begin{aligned} W(n, i, j) &= \frac{\binom{n}{i+1}}{\binom{n}{i}} \frac{\binom{n}{i+2}}{\binom{n}{i+1}} \cdots \frac{\binom{n}{j}}{\binom{n}{j-1}} \left(\binom{j}{i} - \binom{j}{n-i} \right) \\ &= \binom{n}{j} \binom{n}{i}^{-1} \binom{j}{i} - \binom{n}{j} \binom{n}{i}^{-1} \binom{j}{n-i} \\ &= \binom{n-i}{n-j} - \binom{i}{n-j}. \end{aligned}$$

The values of $W(n, i, j)$ is tabulated in the $\lceil n/2 \rceil \times n$ matrix W_n :

$$\begin{pmatrix} \binom{n-0}{n-0} - \binom{0}{n-0} & \binom{n-0}{n-1} - \binom{0}{n-1} & \binom{n-0}{n-2} - \binom{0}{n-2} & \cdots & \binom{n-0}{n-(n-1)} - \binom{0}{n-(n-1)} \\ 0 & \binom{n-1}{n-1} - \binom{1}{n-1} & \binom{n-1}{n-2} - \binom{1}{n-2} & \cdots & \binom{n-1}{n-(n-1)} - \binom{1}{n-(n-1)} \\ 0 & 0 & \binom{n-2}{n-2} - \binom{2}{n-2} & & \binom{n-2}{n-(n-1)} - \binom{2}{n-(n-1)} \\ \vdots & \vdots & & & \vdots \\ 0 & 0 & 0 & & \binom{n-\lceil n/2 \rceil + 1}{n-(n-1)} - \binom{\lceil n/2 \rceil - 1}{n-(n-1)} \end{pmatrix}$$

By removing the leftmost column of W_n we get M_{n-1} . Thus it is sufficient to prove that W_n is totally nonnegative to conclude the same about M_{n-1} . Fomin and Zelevinsky wrote a nice survey on testing total positivity and related questions [3]. We need a result by Lindström [5], and Gessel and Viennot [4].

Lemma 2.2 *If a weighted acyclic directed planar graph has nonnegative real weights, then its weight matrix is totally nonnegative.*

The weight matrix X of a weighted acyclic directed planar graph G , given a set I of sinks and J of sources of G , is a matrix with the rows indexed by I and the columns indexed by J . On the position i, j of X , where $i \in I$ and $j \in J$, is the sum of the weights of all paths from the source i to the sink j .

The graph T_n with its $\lceil n/2 \rceil$ sources and n sinks described earlier gives the weight matrix W_n . All weight are nonnegative, and hence W_n and its submatrix M_{n-1} are totally nonnegative. $\square$

References

- [1] D. André, *Solution directe du problème résolu par M. Bertrand*, C.R. Acad. Sci. Paris **105** (1887), 436–437.
- [2] A. Björner, *A Comparison theorem for f -vectors of simplicial polytopes*, Pure Appl. Math. Q. ([arXiv.org/math/0605336](https://arxiv.org/math/0605336)).
- [3] S. Fomin, A. Zelevinsky, *Total positivity: tests and parametrization*, Math. Intelligencer **22** (2000), no. 1, 23–33.
- [4] I. Gessel, G. Viennot, *Binomial determinants, paths, and hook length formulae*, Adv. in Math. **58** (1985), no. 3, 300–321.
- [5] B. Lindström, *On the vector representations of induced matroids*, Bull. London Math. Soc. **5** (1973), 85–90.

- [6] F.S. Macaulay, *Some properties of enumeration in the theory of modular systems*, Proc. London. Math. Soc, **26** (1927), 531–555.
- [7] S.G. Mohanty, *Lattice path counting and applications*, Probability and Mathematical Statistics. Academic Press, 1979.
- [8] G.M. Ziegler, *Lectures on polytopes*, Graduate Texts in Mathematics, 152. Springer-Verlag, New York, 1995.